



For more information about the Organization of Teratology Information Specialists or to find a service in your area, call (866) 626-6847 or visit us online at: www.OTISpregnancy.org.

E. coli and Pregnancy

This sheet talks about the risks that exposure to E. coli can have during pregnancy. With each pregnancy, all women have a 3% to 5% chance of having a baby with a birth defect. This information should not take the place of medical care and advice from your health care provider.

What is E. coli?

E. coli (*Escherichia coli*) is a bacteria that lives in your colon (gut) and vagina. There are many different types of E. coli bacteria and most are harmless to humans, but some can cause severe illness. Some ways people can get infected with E. coli are:

- Eating contaminated raw and unwashed fruits and vegetables
- Drinking unpasteurized milk and fruit juices
- Eating raw or undercooked meat
- Drinking or swimming in infected water
- Coming into contact with feces from infected farm or petting zoo animals

What are the symptoms of E. coli infection?

Most people will have stomach cramps, slight fever, and diarrhea, but many people have no symptoms. In severe cases, there can be bloody diarrhea, which requires immediate medical care. Rarely people with E. coli infection can develop a form of kidney failure called hemolytic uremic syndrome. This condition is a serious health concern and can lead to kidney damage and death.

How is E. coli infection diagnosed and treated?

If you have symptoms of E. coli infection, a doctor will test a sample of your bowel movement (feces). Most healthy people

recover in a couple of days without the need for antibiotics or over-the-counter medications that stop diarrhea.

Does E. coli infection cause birth defects or pregnancy complications?

There are no reports of E. coli infection causing birth defects in humans. Because diarrhea causes the body to lose a lot of fluids, pregnant women with an E. coli infection can easily become dehydrated. In rare cases, they may start to bleed heavily. There may be a risk for miscarriage or premature delivery with severe E. coli infection. If you think you have an E. coli infection, you should see a doctor right away so that you can be diagnosed and treated if necessary.

If I have an E. coli infection, can I pass it to my unborn baby?

Some types of E. coli that are normally present in the vagina can cause a baby to become infected during delivery. It is not known if E. coli from contaminated food or water can cross the placenta or infect a baby during delivery.

How can I prevent getting an E. coli infection?

- Always wash your hands with soap and water after using or cleaning the

bathroom, changing diapers, handling dirty towels or linens, and touching animals or their environments.

- Always wash your hands after handling raw meat and clean any surface touching raw meat with a disinfectant or bleach and water solution.
- Cook meat thoroughly, especially ground beef, to 160°F. Wash all vegetables and fruits before eating.
- Drink only milk and juice that has been pasteurized.
- Avoid swallowing water when swimming.

I have an E. coli infection. Can I breastfeed my baby?

Yes. The E. coli bacteria do not get into the breast milk, so it is okay to continue breastfeeding. There are important immune factors in breast milk that can help protect your baby from infections. Having diarrhea and other symptoms of E. coli infection may cause a decrease in your milk supply, so drink plenty of fluids. Make sure that you wash your hands before you hold or breastfeed your baby. If your baby gets diarrhea or other symptoms of E. coli infection, contact your pediatrician immediately.

July 2010.
Copyright by OTIS.
Reproduced by permission.



Selected References:

Bower J. 1999. Foodborne diseases: shiga toxin producing E. coli (STEC). *Pediatr Infect Dis J*. 18(10):909-910.

Chalupka S. 2005. Tainted water on tap: what to tell patients about preventing illness from drinking water. *Am J Nurs*. 105(11):40-52.

Committee on Infectious Diseases, American Academy of Pediatrics. 2003. Red Book: 2003 Report of the Committee on Infectious Diseases. 26th ed. Elk Grove Village (IL): American Academy of Pediatrics.

Gaither K, Ardite A, and Mason TC. 2005. Pregnancy complication by emphysematous pyonephrosis. *J Natl Med Assoc*. 97(10):1411-3.

Jones B, et al. 2004. *Escherichia coli*: a growing problem in early onset neonatal sepsis. *Aust N Z J Obstet Gynaecol*. 44:558-561.

Palmeira P, et al. 2005. Colostrum from healthy Brazilian women inhibits adhesion and contains IgA antibodies reactive with Shiga toxin-producing *Escherichia coli*. *Eur J Pediatr*. 164(1):37-43.

Pickering L, Guerrant RL and Cleary TG. 2001. Nature of the Organism: Microorganisms Responsible for Neonatal Diarrhea. In: Remington JS and Klein JO, editors. Infectious Diseases of the Fetus and Newborn Infant. 5th Ed. Philadelphia: W.B. Sanders Co. p.1251-1265.

Schraq SJ, et al. 2006. Risk factors for invasive, early-onset *Escherichia coli* infections in the era of widespread intrapartum antibiotic use. *Pediatrics*. 118(2):570-6.

Watt S, et al. 2003. *Escherichia coli* strains from pregnant women and neonates: intraspecies genetic distribution and prevalence of virulence factors. *J Clin Microbiol*. 41(5):1929-1935.

*If you have questions about the information on this fact sheet or other exposures during pregnancy, call **OTIS** at **1-866-626-6847**.*