



ORGANIZATION  
OF TERATOLOGY  
INFORMATION  
SERVICES

For more information regarding OTIS or a Teratology Information Service in your area, call OTIS Information at (866) 626-6847 or visit us online at: [www.OTISpregnancy.org](http://www.OTISpregnancy.org).

## **Lymphocytic Choriomeningitis Virus (LCMV) and Pregnancy**

The information below will help you determine if your exposure to LCMV represents an increased fetal risk. With every pregnancy, all women have a 3 to 5 percent chance to have a baby with a birth defect.

### ***What is Lymphocytic Choriomeningitis Virus (LCMV)?***

LCMV is a virus that causes symptoms similar to other viral infections, including fever, muscle aches, fatigue and gastrointestinal problems. LCMV is not thought to be a common virus in humans. The Centers for Disease Control estimates that 2% to 10% of adults have had an LCMV infection. Rodents such as mice and hamsters carry the virus. Rodents kept as pets, in laboratories and in the wild have all been found to carry LCMV.

### ***How could I get LCMV?***

Infected rodents can transfer the infection to other rodents and humans. Physical contact or accidental ingestion of rodents' saliva, urine, feces or blood may cause an LCMV infection. Sweeping rodent droppings may cause the virus to become airborne and increase the chances of getting an infection. Human infection with this virus is more common in the fall when rodents move into homes.

### ***How can I prevent getting an LCMV infection?***

The best way to prevent LCMV is to minimize or avoid direct physical contact with rodents and their body fluids. A

pregnant woman should ask someone else to change the litter of pet rodents.

When handling or cleaning rodents or their cages use gloves and wash hands thoroughly after contact. When cleaning litter, use a diluted bleach solution on visible droppings and surrounding areas. In areas that are heavily populated with rodents (such as barns, storage sheds, labs) adequate ventilation minimizes exposure. Wetting the area may also prevent the virus from becoming airborne.

### ***How will I know if I have LCMV?***

In humans, LCMV infection usually causes mild illness such as fever, fatigue, loss of appetite, muscle aches, headache, nausea and vomiting. A small number of individuals may become quite ill and develop meningitis or encephalitis (inflammation of the brain or spinal cord). Other individuals may not have any symptoms. Your doctor may be able to order a blood test to detect this infection.

### ***If I'm in my first trimester, will an LCMV infection cause a miscarriage?***

During pregnancy, a woman who gets an LCMV infection may have an increased chance for a miscarriage. The data is very limited and the exact risks for miscarriage are unknown.

### ***How does LCMV affect the developing baby?***

A small number of children (49 total) have been reported worldwide with congenital LCMV (present at birth). Infants described as having congenital LCMV may have abnormalities in brain and eye development, as well as learning difficulties. The most common birth defects are hydrocephalus (fluid in the brain) and chorioretinitis (eye problems which can lead to vision loss).

It is unknown how many women have had LCMV during pregnancy and had healthy babies. Because of the limited amount of information about LCMV in humans, the exact risks for LCMV related birth defects are unknown.

### ***If I get LCMV when I'm pregnant, how is it treated?***

No special guidelines exist for treatment of LCMV during pregnancy; however, your doctor may prescribe medication if a blood test shows a recent LCMV infection. As with any viral illness, appropriate treatment of symptoms (such as fever) due to the infection is recommended.

### ***What if I have LCMV while I'm breastfeeding?***

No information is available about LCMV and breastfeeding.



ORGANIZATION  
OF TERATOLOGY  
INFORMATION  
SERVICES

**July 2003.**  
**Copyright by OTIS.**  
**Reproduced by permission.**

### **References**

Barton LL, Mets MB. 2001. Congenital Lymphocytic Choriomeningitis Virus Infection: Decade of Rediscovery. *CID* 33:370-4.

Barton LL, Peters CJ, Ksiazek TG. 1995. Lymphocytic Choriomeningitis Virus: An Unrecognized Teratogenic Pathogen. *Emerging Infectious Diseases* 1(4): 152-3.

Braude AI, Davis CE, Fierer J. eds 1986. *Infectious Diseases and Medical Microbiology*, 2<sup>nd</sup> ed., W.B. Saunders Co., 1076-1078, 60.

Childs JE, Glass GE, Korch GW, Ksiazek TG, Leduc JW. 1992. Lymphocytic Choriomeningitis Virus Infection and House Mouse (*Mus Musculus*) Distribution in Urban Baltimore. *Am J Trop Med Hyg* 47(1): 27-34.

Hirsch MS, Moellering RC, Pope HG, Poskanzer DC. 1974. Lymphocytic-Choriomeningitis-Virus Infection Traced to a Pet Hamster. *NEJM*. 291(12): 610-611.

Jenson K. "University of Arizona Study Published in *Clinical Infectious Diseases* Warns Pregnant Women about the Risk of Rodents," press release, August 9, 2001.

Komrower GM, Williams BL, Stones PB. 1955. Lymphocytic Choriomeningitis in the Newborn: Probable Transplacental Infection. *Lancet* 1:697-8.

Mandell, Douglas, Bennett's Principle and Practice of Infectious Diseases, 5<sup>th</sup> ed., Vol. 2, Philadelphia: Churchill Livingstone, 2000, pp.1855-57.

Traube E. 1973. LCM Virus Research, Retrospect and Prospects Lymphocytic Choriomeningitis Virus and Other Arenaviruses, Lehmann-Grube, F. ed., Springer-Verlag: Berlin, pp.3-10.

Center for Disease Control. Lymphocytic Choriomeningitis. Retrieved June 2003, from: <http://www.cdc.gov/ncidod/dvrd/spb/mnpages/di/pages/lcmv.htm>