



For more information regarding OTIS or a Teratology Information Service in your area, call OTIS Information at (866) 626-6847 or visit us online at: [www.OTISpregnancy.org](http://www.OTISpregnancy.org).

## Tetracycline and Pregnancy

The information below will help you determine if your prenatal exposure to tetracycline represents an increased fetal risk. With every pregnancy, all women have a 3 to 5 percent chance to have a baby with a birth defect.

### ***What is tetracycline?***

Tetracycline is an antibiotic used to treat conditions including acne and respiratory infections. It belongs to a group of antibiotics that includes minocycline, oxytetracycline, doxycycline, and other related antibiotics. In general, the concerns stated in this fact sheet apply to tetracycline and these related antibiotics. Tetracycline is usually taken orally.

### ***I am taking tetracycline during the first month of my pregnancy. Is there a risk for birth defects?***

Researchers have studied the outcome of pregnancies when women took tetracycline during the first trimester (the first 12 weeks), which is the period when all the body organs are forming. These studies have not shown an increased risk for major birth defects in children exposed to tetracycline in the first trimester of pregnancy.

There may be a small increased risk for minor birth defects such as an inguinal hernia, but the number of reports is too small to be sure this risk exists. Once you know that you are pregnant, it is a good idea to talk to your health care provider about continuing the use of tetracycline for the remainder of your pregnancy.

### ***Is there a risk for other problems if I take tetracycline at any time during pregnancy?***

Yes. If you take tetracycline after the fourth month of pregnancy, there is a risk for

discoloration (graying) of the "baby" teeth. The discoloration of the baby teeth is due to calcification (hardening) of the teeth, which starts at about four months. This problem appears to be only cosmetic, and does not appear to affect the development of the enamel or the chance of getting cavities. We do not know what percentage of exposed babies will have discoloration of the teeth. For this reason, the use of tetracycline should be avoided after four months of pregnancy unless there is a special condition requiring the doctor to prescribe this type of antibiotic.

### ***I have heard that tetracycline also affects the bones of developing babies, in addition to the teeth. Is this true?***

Tetracycline appears to affect the calcification (hardening) of the bones and teeth, and it also appears to cause reduced growth of some bones while the baby is being exposed to the medication. Yet, while the tooth discoloration is permanent, the bone growth seems to return to normal after the tetracycline exposure ends.

### ***I am 14 weeks pregnant and I am taking tetracycline. If I stop taking it now, how long will tetracycline stay in my body?***

It takes less than 5 days for tetracycline to be cleared from the body of most people. If you stop taking the medication now, your baby will not be exposed to tetracycline after one week. As with any medication, you should speak with your health care provider before you stop taking tetracycline.

***What if I am taking one of the other medications in this group such as minocycline, oxytetracycline, or doxycycline instead of tetracycline? Does that still put my baby at risk?***

Since these antibiotics are similar to tetracycline, it is possible that they could also affect a baby's teeth and bones in the same way if taken in the second and third trimesters of pregnancy. There are some reports of babies with dental discoloration after similar exposures, and for that reason, it is best to use the same precautions with these related medications.

***I am pregnant and I often get upper respiratory infections for which my doctor prescribes tetracycline. If I get one of these infections after I am five months pregnant, what should I do?***

Talk to your doctor about your concerns regarding tetracycline. Your doctor may consider a different medication to treat your infection.

***I am breast feeding and thinking about taking tetracycline. Can it still affect my baby?***

In a baby, the bones and teeth continue to harden until one year of age. Therefore, if a baby is exposed to tetracycline in breast milk, theoretically it could cause tooth discoloration and delayed bone growth. Yet, since only very low levels of tetracycline are secreted into the breast milk, no problems have been reported in nursing babies.

In fact, the American Academy of Pediatrics has approved tetracycline for use during breastfeeding. Also, the WHO Working Group on Human Lactation states that when the antibiotic is used for seven to 10 days while nursing, the risk to the infant appears to be low. If you have any concerns about breastfeeding while taking these types of antibiotics, talk to your health care provider.

November 2000  
Copyright by OTIS,  
Reproduced by permission.



**References:**

- Briggs et al. Drugs in Pregnancy and Lactation, 5<sup>th</sup> Ed. Williams and Wilkins (1998).
- Cohlan, SQ et al. (1961) "Effect of tetracycline on bone growth in the premature infant." *Antimicrobial Agents Chemotherapy*. 1961: 340-7.
- Committee on Drugs, American Academy of Pediatrics (1994). "The transfer of drugs and other chemicals into human breast milk." *Pediatrics*. 93: 137-150.
- Egerman, et al. (1992). "The tetracyclines." *Obstet Gynecol Clin NA*. 19(3): 551-61.
- Genot, MT et al. (1970). "Effect of administration of tetracycline on teeth." *J Oral Med*. 25: 75-9.
- Kline et al. (1964). "Transplacental effect of tetracycline on teeth." *JAMA*. 188: 178.
- Kutscher et al. (1966). "Discoloration of deciduous teeth induced by administration of tetracycline antepartum." *Am J Obstet Gynecol*. 96: 291-2.
- The WHO Working Group, Bennet, PN (Ed): Drugs and Human Lactation. Elsevier, New York, (1988).
- Gilman, AG et al. (Eds). Goodman and Gilman's The Pharmacological Basis of Therapeutics, 7<sup>th</sup> Ed., Macmillan, New York, (1985).
- O'Rahilly, R, and Mueller, F. Human Embryology & Teratology. Wiley-Liss, New York, (1992).